

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
13 January 2005 (13.01.2005)

PCT

(10) International Publication Number
WO 2005/003166 A1

(51) International Patent Classification⁷: **C07K 14/705**,
5/09, A61K 38/17, 38/06, 49/00, G01N 33/58, 33/68

(21) International Application Number:
PCT/NO2004/000208

(22) International Filing Date: 7 July 2004 (07.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
20033115 8 July 2003 (08.07.2003) NO

(71) Applicant (for all designated States except US): **AMERSHAM HEALTH AS** [NO/NO]; Intellectual Property Dept., P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CUTHBERTSON, Alan** [GB/NO]; Amersham Health AS, P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO). **INDREVOLL, Bård** [NO/NO]; Amersham Health AS, P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO). **SOLBAKKEN, Magne** [NO/NO]; Amersham Health AS, P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO).

(74) Agents: **WULFF, Marianne, Weiby et al.**; Amersham Health AS, P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FLUORESCCEIN-LABELLED PEPTIDES

(57) Abstract: The invention relates to new peptide-based compounds and their use in diagnostic optical imaging. More specifically the invention relates to the use of such peptide-based compounds as targeting vectors that bind to receptors associated with angiogenesis. The compounds are labeled with fluorescein and may be used as contrast agents in optical imaging in diagnosis of angiogenesis-related diseases.



WO 2005/003166 A1

FLUORESCCEIN-LABELLED PEPTIDES

Field of the invention

This invention relates to new peptide-based compounds and
5 their use in diagnostic optical imaging or for the
treatment of disease. More specifically the invention
relates to the use of such peptide-based compounds as
targeting vectors that bind to receptors associated with
angiogenesis. The compounds may be used as contrast
10 agents in diagnosis of angiogenesis- related diseases.

Background of invention

Generally, new blood vessels can be formed by two
different mechanisms: vasculogenesis or angiogenesis.
15 Angiogenesis is the formation of new blood vessels by
branching from existing vessels. The primary stimulus
for this process may be inadequate supply of nutrients
and oxygen (hypoxia) to cells in a tissue. The cells may
respond by secreting angiogenic factors, of which there
20 are many; one example, which is frequently referred to,
is vascular endothelial growth factor (VEGF). These
factors initiate the secretion of proteolytic enzymes
that break down the proteins of the basement membrane, as
well as inhibitors that limit the action of these
25 potentially harmful enzymes. The other prominent effect
of angiogenic factors is to cause endothelial cells to
migrate and divide. Endothelial cells that are attached
to the basement membrane, which forms a continuous sheet
around blood vessels on the contraluminal side, do not
30 undergo mitosis. The combined effect of loss of
attachment and signals from the receptors for angiogenic
factors is to cause the endothelial cells to move,
multiply, and rearrange themselves, and finally to
synthesise a basement membrane around the new vessels.

Angiogenesis is prominent in the growth and remodelling of tissues, including wound healing and inflammatory processes. Inhibition of angiogenesis is also considered
5 to be a promising strategy for antitumour therapy. The transformations accompanying angiogenesis are also very promising for diagnosis, an obvious example being malignant disease, but the concept also shows great
10 promise in inflammation and a variety of inflammation-related diseases, including atherosclerosis, the macrophages of early atherosclerotic lesions being potential sources of angiogenic factors. These factors are also involved in re-vascularisation of infarcted parts of the myocardium, which occurs if a stenosis is
15 released within a short time.

Further examples of undesired conditions that are associated with neovascularization or angiogenesis, the development or proliferation of new blood vessels are
20 shown below. Reference is also made in this regard to WO 98/47541.

Diseases and indications associated with angiogenesis are e.g. different forms of cancer and metastasis, e.g.
25 breast, skin, colorectal, pancreatic, prostate, lung or ovarian cancer.

Other diseases and indications are inflammation (e.g. chronic), atherosclerosis, rheumatoid arthritis and
30 gingivitis.

Further diseases and indications associated with angiogenesis are arteriovenous malformations, astrocytomas, choriocarcinomas, glioblastomas, gliomas,

hemangiomas (childhood, capillary), hepatomas, hyperplastic endometrium, ischemic myocardium, endometriosis, Kaposi sarcoma, melanoma, neuroblastomas, occluding peripheral artery disease, osteoarthritis, psoriasis, retinopathy (diabetic, proliferative), scleroderma, seminomas and ulcerative colitis.

Angiogenesis involves receptors that are unique to endothelial cells and surrounding tissues. These markers include growth factor receptors such as VEGF and the Integrin family of receptors. Immunohistochemical studies have demonstrated that a variety of integrins, perhaps most importantly the α_v class, are expressed on the apical surface of blood vessels [Conforti, G., et al. (1992) Blood 80: 37-446] and are available for targeting by circulating ligands [Pasqualini, R., et al. (1997) Nature Biotechnology 15: 542-546]. The $\alpha_5\beta_1$ is also an important integrin in promoting the assembly of fibronectin matrix and initiating cell attachment to fibronectin. It also plays a crucial role in cell migration.

The integrin $\alpha_v\beta_3$ is one of the receptors that is known to be associated with angiogenesis. Stimulated endothelial cells appear to rely on this receptor for survival during a critical period of the angiogenic process, as antagonists of the $\alpha_v\beta_3$ integrin receptor/ligand interaction induce apoptosis and inhibit blood vessel growth.

Integrins are heterodimeric molecules in which the α - and β -subunits penetrate the cell-membrane lipid bilayer. The α -subunit has four Ca^{2+} binding domains on its extracellular chain, and the β -subunit has a number of extracellular cysteine-rich domains.

Many ligands (eg. fibronectin) involved in cell adhesion contain the tripeptide sequence arginine-glycine-aspartic acid (RGD). The RGD sequence appears to act as a primary
5 recognition site between the ligands presenting this sequence and receptors on the surface of cells. It is generally believed that secondary interactions between the ligand and receptor enhance the specificity of the interaction. These secondary interactions might take
10 place between moieties of the ligand and receptor that are immediately adjacent to the RGD sequence or at sites that are distant from the RGD sequence.

RGD peptides are known to bind to a range of integrin
15 receptors and have the potential to regulate a number of cellular events of significant application in the clinical setting. Perhaps the most widely studied effect of RGD peptides and mimetics thereof relate to their use as anti-thrombotic agents where they target the platelet
20 integrin GpIIbIIIa.

Inhibition of angiogenesis in tissues by administration of either an $\alpha v\beta 3$ or $\alpha v\beta 5$ antagonist has been described in for example WO 97/06791 and WO 95/25543 using either
25 antibodies or RGD containing peptides. EP 578083 describes a series of mono-cyclic RGD containing peptides.

Cyclic RGD peptides containing multiple bridges have also
30 been described in WO 98/54347 and WO 95/14714.

Further examples of RGD comprising peptide-based compounds are found in WO01/77145, WO02/26776 and WO03/006491.

There is a clinical need to develop more specific non-invasive imaging techniques for angiogenesis-related diseases. Such imaging techniques will have a central
5 role in the evaluation of novel anti-angiogenic therapies. Being able to assess the actual level of angiogenesis will be of clinical benefit in diagnosing angiogenesis-related diseases at an early stage. It has now surprisingly been found that optical imaging may be
10 used to assess the level of angiogenesis, and the invention provides new optical imaging contrast agents for this purpose.

Summary of the invention

15 In view of the needs of the art the present invention provides peptide-based compounds labeled with fluorescein for use as contrast agents in optical imaging or for therapeutic treatment. The efficient targeting and imaging of integrin receptors associated with
20 angiogenesis in vivo demands a selective, high affinity RGD based vector that is chemically robust and stable. Furthermore, the route of excretion is an important factor when designing imaging agents in order to reduce problems with background noise. These stringent
25 conditions are met by the fluorescein-labelled peptide compounds described in the present invention.

Detailed description of the invention

Viewed from one aspect the invention provides new
30 peptide-based compounds as defined in the claims. These compounds have affinity for integrin receptors, e.g. affinity for the integrin $\alpha v \beta 3$, and are labeled with a fluorescein dye reporter.


The compounds, or physiologically acceptable salts thereof, comprise a peptidic vector and at least one fluorescein dye, wherein the peptidic vector comprises the amino acid sequence X_3 -G-D and wherein the peptidic vector and the fluorescein dyes are coupled, preferably by covalent bonds. X_3 represents arginine, N-methylarginine or an arginine mimetic, G represents glycine and D represents aspartic acid. The peptidic vector has affinity for integrin receptors, such as the $\alpha v \beta 3$ receptors.

The compounds of the invention comprise the amino acid sequence X_3 -G-D having affinity for the integrin receptors. The compound preferably comprises further amino acids, and optional other moieties, wherein the X_3 -G-D sequence is the binding seat of the peptidic vector which function as a vector binding to an integrin type receptor.

The compound of the invention can be constrained for example by formation of one or more cyclising bridges in the peptidic vector part. A monocyclic peptide compound can be obtained by formation of a disulfide bond or a thioether bond between amino acids. A peptide-based compound including one cyclising bridge is more specific towards $\alpha v \beta 3$, and is more preferred, than a linear peptide. The compounds of the invention preferably comprise two cyclising bridges between different amino acids of the compounds. The term "cyclising bridges" refers to any combination of amino acids or with amino acids and $-(CH_2)_n-$ or $-(CH_2)_n-C_6H_4-$ groups with functional groups which allows for the introduction of a bridge. n represents a positive integer from 1 to 10. Some preferred examples are disulphides, disulphide mimetics such as the $-(CH_2)_4-$ carba bridge, thioacetal, thioether bridges (cystathione or lanthionine) and bridges containing esters and ethers. Preferably, one bridge

forms a disulphide bond and a second bridge comprises a thioether (sulphide) bond.

In a further embodiment the compounds of the invention
5 are identified by formula(I)



comprising two cyclising bridges,
wherein,

10 X₃, G and D are as previously defined; and

R_a represents -(CH₂)_n- or -(CH₂)_n-C₆H₄- forming a bridge to either of X₂, X₄ or X₆, wherein

n represents a positive integer from 1 to 10; and

15

X₁ represents a bond or 1, 2, 3, 4 or 5 amino acid residues, wherein one amino acid residue is optionally functionalised with a spacer moiety, and preferably said amino acid residue possesses a functional side-chain such
20 as an acid or amine group preferably selected from aspartic or glutamic acid, lysine, homolysine, diaminoalcylic acid or diaminopropionic acid; and


X₂ and X₄ represent independently amino acids residues
25 capable of forming a cyclising bridge, such as cysteine or homocysteine residues forming disulphide or thioether bonds, or other amino acid residues capable of forming a cyclising bridge such as aspartic acid and lysine, preferably X₂ and X₄ represent residues of cysteine or
30 homocysteine; and

X₅ represents a hydrophobic amino acid or derivatives thereof, and preferably represents a tyrosine, a phenylalanine, a 3-iodo-tyrosine or a naphthylalanine

residue, and more preferably a phenylalanine or a 3-iodo-tyrosine residue; and

X_6 represents an amino acid residue capable of forming a cyclising bridge, preferably a thiol-containing amino-acid residue, preferably a cysteine or a homocysteine residue; and

X_7 represents a spacer or biomodifier moiety or is absent, and is preferably based on a monodisperse polyethylene glycol (PEG) building block comprising 1 to 10 units of said building block, said biomodifier having the function of modifying the pharmacokinetics and blood clearance rates of said agents. In addition X_7 may also represent 1 to 10 amino acid residues preferably comprising glycine, lysine, aspartic acid or serine. More preferably, X_7 represent a spacer or biomodifier comprising both amino acid residues and a PEG-like structure, such as 1-10 amino acid residues in combination with a PEG-like structure, preferably a bis aminoethyl ethylene glycol glycine combination. In a preferred embodiment X_7 represents a unit comprised of the monodisperse PEG-like structure, 17-amino-5-oxo-6-aza-3,9,12,15-tetraoxaheptadecanoic acid of formula (II),



(II)

wherein m equals an integer from 1 to 10 and where the C-terminal end is an amide or acid moiety. It is found that the biomodifier, X_7 , modifies the pharmacokinetics and blood clearance rates of the compounds. The biomodifier

effects less uptake of the compounds in tissue i.e. muscle, liver etc. thus giving a better diagnostic image due to less background interference. The secretion is mainly through the kidneys due to a further advantage of
5 the biomodifier; and

the compound further comprising at least one group Z, representing a fluorescein group, linked to one or more of X₁, X₆ or X₇, optionally via a spacer group.

10

The peptide-based compound comprises a peptidic vector defined by the amino sequence formed by X₁, X₂, X₃, G, D, X₄, X₅ and X₆ of Formula I and this peptide constitute a targeting vector having affinity for integrin receptors
15 associated with angiogenesis.


Depending of the placement of the cyclising bridges the compounds will comprise "discrete", "nested" or "interlocking" configurations. Preferably the two bridges
20 in each compounds are:

Between R_a and X₆, and between X₂ and X₄ (forming a nested configuration);

Between R_a and X₂, and X₄ and X₆ (discrete configuration);


25 Between R_a and X₄, and X₂ and X₆ (forming an interlocking configuration).

In a further embodiment the compounds of the invention are identified by either of the formulas below:




(III)

5



(IV)

10



(V)

15

wherein R_a , X_1 , X_3 , G , D , X_5 , and X_7 are as defined for formula I; and wherein

20 X_2 , X_4 and X_6 represent amino acid residues capable of forming a disulphide or a thioether bond, such as cysteines or homocysteines;

W_1 is a spacer moiety or is absent, and is preferentially
 25 derived from glutaric and/or succinic acid and/or a

polyethyleneglycol based unit, e.g. including a moiety as given for formula II, linking the fluorescein dye to the peptide. Other representative spacer (W_1) elements include structural-type polysaccharides, storage-type polysaccharides, polyamino acids and methyl and ethyl esters thereof, and polypeptides, oligosaccharides and oligonucleotides, which may or may not contain enzyme cleavage sites. The role of the spacer moiety W_1 is to distance the relatively bulky dye from the receptor binding domain of the peptide component;

h is a positive integer 1 or 2;

and wherein at least one of the Z groups is present representing a fluorescein dye.

The compounds preferably include only one Z -group.

The fluorescein dye, represented by Z may be linked to X_1 , W_1 , X_6 or X_7 , preferably by amide bond formation.

In a preferred aspect the compounds of formula III-V, or the physiologically acceptable salts thereof, have the following characteristics:

R_a preferably represents $-(CH_2)-$.

Further, X_1 represents an amino acid residue with a functional side-chain such as an acid or amine group, the amino acid preferably being selected from aspartic or glutamic acid, lysine, homolysine, diaminoalicyclic acid or diaminopropionic acid, more preferably aspartic acid or lysine.

X₂, X₄ and X₆ preferably independently represent a cysteine or a homocysteine residue.

X₃ preferably represents arginine.

5

X₅ preferably represents tyrosine, phenylalanine, 3-iodo-tyrosine or naphthylalanine, and more preferably phenylalanine or 3-iodo-tyrosine.

10 X₇ and W₁ are defined as for formula I and II-V respectively. Preferably X₇ comprises 1-10 units of a monodisperse PEG building block or is absent, and W₁ is preferably absent.


15 Z represents a fluorescein dye or is absent, such that the compound comprises at least one fluorescein dye.

In a preferred aspect the compounds are of formula III (nested) or physiologically acceptable salts thereof, and
20 more preferably they have the characteristics given in the preferred aspect described above.

The compounds of formula III-V comprise at least two bridges, wherein one bridge forms a disulphide bond and
25 the second bridge comprises a thioether (sulphide) bond. The preferred compounds, of formula III, have the bridges placed such the bridges fold the peptide moiety into a 'nested' configuration. The compounds of this embodiment of the invention thus have a maximum of one disulphide
30 bridge per molecule moiety. Compounds defined by the present invention are surprisingly stable in vivo and under the conditions employed during labeling with fluorescein.


The reporter Z comprises a fluorescein dye, i.e. fluorescein or derivatives of fluorescein. Fluorescein is a yellow dye which glows in visible light. Fluorescein is typically excited by the 488 nm line of an argon laser, and emission is collected at 530 nm. Fluorescein and its derivatives have relatively high absorptivity, excellent fluorescence quantum yield and good water solubility. The extensive use in the past makes them very well characterised, the availability is very good and they have a general low cost per mg.

Formula (VI) gives fluorescein, showing the numbering of the different positions.



(VI) Fluorescein


Relevant derivatives of fluorescein are e.g. the carboxyfluoresceins, such as the 5-carboxyfluorescein, shown by formula VII, or the 6-carboxyfluorescein.



5

(VII) 5-carboxyfluorescein

Another relevant fluorescein derivative is fluorescein isothiocyanate given by formula VIII:




10

(VIII) fluorescein isothiocyanate

Even more preferred is to use an activated carboxylate of
15 fluorescein, such as an N-hydroxysuccinimide (NHS) ester,
called fluorescein NHS ester, shown in formula IX.
Succinimidyl esters are excellent reagents since the
amide products formed are very stable.

20

15



(IX) Fluorescein NHS ester

5

The fluorescein dye, represented by Z, may be linked to X₁, X₆ or X₇ of formula I, or optionally via the spacer W₁ as given in formulas III-V. Preferably, fluorescein is coupled to the peptidic vector by amide bond formation with a suitable amino group of the peptide. Active esters of fluorescein such as the NHS ester (Pierce Catalog 46100) are considered particularly useful when synthesising the compounds. Preferred coupling sites for the fluorescein are position 5 and 6.

15


The amino acids may preferably represent a naturally occurring amino acid. In most cases, it is preferred that the amino acids in the peptide are all in the L-form. However, in some embodiments of the invention one, two, three or more of the amino acids in the peptide are preferably in the D-form. The inclusion of such D-form amino acids can have a significant effect on the serum stability of the compound.

Some of the compounds of the invention are high affinity RGD based vectors. As used herein the term 'high affinity RGD based vector' refers to compounds that have

a K_i of < 10 nM and preferably < 5 nM, in a competitive binding assay for $\alpha v \beta 3$ integrin and where the K_i value was determined by competition with the known high affinity ligand echistatin. Methods for carrying out such competition assays are well known in the art.


Some preferred compounds of the invention are illustrated below. Compounds A and B comprise fluorescein conjugated to an RGD-containing peptide (Lys-Cys-Arg-Gly-Asp-Cys-Phe-Cys by amide bond formation, both giving a "nested" configuration.

Compound A:



15


Compound B:




Other examples of compounds of the invention are illustrated by compounds C and D below.

Compound C:

- 5 The compound comprises an RGD peptide (Lys-Asp-Cys-Arg-Gly-Asp-Cys-Phe-Cys-Gly) linked to fluorescein, forming a "discrete" configuration.



- 10 Compound D:
The compound comprises an RGD-peptide (Lys-Cys-Arg-Gly-Asp-Cys-Phe-Cys) linked to fluorescein, forming an "interlocking" configuration.



The new compounds of the invention may be used as contrast agents in optical imaging or for treatment of diseases. Use of the compounds as described for treatment or as contrast agents in optical imaging is hence an
5 embodiment of the invention. A preferred embodiment of the invention is compounds as described for use as optical imaging contrast agent, particularly for use in optical imaging of and diagnosing of angiogenesis-related diseases.

10

The compounds of this invention are useful as imaging agents in the detection of angiogenesis in both humans and animals. The products may also have utility in pre-clinical animal models and allow monitoring of
15 therapeutic efficacy of new drugs within pharmaceutical research, e.g. in oncology.

20

The present invention also provides a pharmaceutical composition comprising an effective amount, e.g. an
amount effective for enhancing image contrast in in vivo imaging of a compound of the invention, or a salt thereof, together with one or more pharmaceutically acceptable adjuvants, excipients or diluents.

25

The invention further provides a pharmaceutical composition for treatment of a disease comprising an effective amount of a compound of the invention, or a salt thereof, together with one or more pharmaceutically acceptable adjuvants, excipients or diluents.

30

Viewed from a further aspect the invention provides the use of a compound of the invention for the manufacture of an optical imaging contrast medium for use in a method of diagnosis involving administration of said contrast

medium to a human or animal body and generation of an image of at least part of said body.

5 Use of the compounds in the manufacture of therapeutic compositions (medicament) and in methods of therapeutic or prophylactic treatment, preferably treatment of angiogenesis-related diseases, of the human or animal body are thus considered to represent further aspects of the invention.

10

Viewed from a still further aspect the invention provides a method of generating an image of a human or animal body by optical imaging involving administering a contrast agent to said body, e.g. into the vascular system and
15 generating an image of at least a part of said body, to which said contrast agent has distributed, wherein as said contrast agent is used a compound as described.

Viewed from a still further aspect the invention provides
20 a method of generating enhanced images of a human or animal body by optical imaging previously administered with a contrast agent composition comprising a compound as defined, which method comprises generating an image of at least part of said body.

25

Viewed from a further aspect the invention provides a method of monitoring the effect of treatment of a human or animal body with a drug to combat a condition associated with angiogenesis, said method involving
30 administering to said body an agent of the invention and detecting the uptake of said agent by cell receptors, preferably endothelial cell receptors and in particular $\alpha v\beta 3$ receptors, said administration and detection optionally but preferably being effected repeatedly, e.g.

before, during and after treatment with said drug. Said detection comprising an optical imaging technique.

Since fluorescein as well as the fluorescein-peptide
5 compound of the present invention emits in the visual range of the spectrum, conventional ophthalmoscopy equipment may be used for acquiring the images. Furthermore, in-vivo confocal microscopy may be applied. Recently developed time-domain and frequency-domain
10 imaging techniques may potentially also be used, taking advantage of additional characteristics of the fluorophore, such as lifetime.


The compounds of the present invention can be synthesized
15 using all the known methods of chemical synthesis but particularly useful is the solid-phase methodology of Merrifield employing an automated peptide synthesizer (J. Am. Chem. Soc., 85: 2149 (1964)). In addition, coupling of fluorescein, such as a fluorescein active ester, can
20 also be carried out automatically yielding an amide bond between the peptidic vector and the fluorescein group. The peptidic vector and the peptide compound may be purified using high performance liquid chromatography (HPLC) and characterized by mass spectrometry and
25 analytical HPLC before testing in the in vitro screen.

The present invention will now be further illustrated by way of the following non-limiting examples.

ExamplesExample 1:

Synthesis of disulphide [Cys²⁻⁶] thioether cyclo[CH₂CO-
 5 Lys (fluorescein)-Cys²-Arg-Gly-Asp-Cys⁶-Phe-Cys]-PEG-NH₂

1 a) Synthesis of 17-(Fmoc-amino)-5-oxo-6-aza-3,9,12,15-
tetraoxaheptadecanoic acid



10

This building block is coupled to the solid-phase using Fmoc chemistry. The coupled form of this building block will be referred to in short as PEG.

15 1,11-Diazido-3,6,9-trioxaundecane

A solution of dry tetraethylene glycol (19.4 g, 0.100 mol) and methanesulphonyl chloride (25.2 g, 0.220 mol) in dry THF (100 ml) was kept under argon and cooled to 0 °C in an ice/water bath. To the flask was added a solution of triethylamine (22.6 g, 0.220 mol) in dry THF (25 ml) dropwise over 45 min. After 1 hr the cooling bath was removed and stirring was continued for 4 hrs. Water (60 ml) was added. To the mixture was added sodium hydrogencarbonate (6 g, to pH 8) and sodium azide (14.3 g, 0.220 mmol), in that order. THF was removed by distillation and the aqueous solution was refluxed for 24 h (two layers formed). The mixture was cooled and ether (100 ml) was added. The aqueous phase was saturated with sodium chloride. The phases were separated and the aqueous phase was extracted with ether (4 x 50 ml).

Combined organic phases were washed with brine (2 x 50 ml) and dried (MgSO₄). Filtration and concentration gave 22.1 g (91%) of yellow oil. The product was used in the next step without further purification.

5

11-Azido-3,6,9-trioxaundecanamine

To a mechanically, vigorously stirred suspension of 1,11-diazido-3,6,9-trioxaundecane (20.8 g, 0.085 mol) in 5% hydrochloric acid (200 ml) was added a solution of
10 triphenylphosphine (19.9 g, 0.073 mol) in ether (150 ml) over 3 hrs at room temperature. The reaction mixture was stirred for additional 24 hrs. The phases were separated and the aqueous phase was extracted with dichloromethane (3 x 40 ml). The aqueous phase was cooled in an ice/water
15 bath and pH was adjusted to ca 12 by addition of KOH. The product was extracted into dichloromethane (5 x 50 ml). Combined organic phases were dried (MgSO₄). Filtration and evaporation gave 14.0 g (88%) of yellow oil. Analysis by MALDI-TOF mass spectroscopy (matrix: α -cyano-4-
20 hydroxycinnamic acid) gave a M+H peak at 219 as expected. Further characterisation using ¹H (500 MHz) and ¹³C (125 MHz) NMR spectroscopy verified the structure.

17-Azido-5-oxo-6-aza-3,9,12,15-tetraoxaheptadecanoic acid

25

To a solution of 11-azido-3,6,9-trioxaundecanamine (10.9 g, 50.0 mmol) in dichloromethane (100 ml) was added diglycolic anhydride (6.38 g, 55.0 mmol). The reaction mixture was stirred overnight. HPLC analysis (column
30 Vydac 218TP54; solvents: A = water/0.1% TFA and B = acetonitrile/0.1% TFA; gradient 4-16% B over 20 min; flow 1.0 ml/min; UV detection at 214 and 284 nm), showed complete conversion of starting material to a product with retention time 18.3 min. The solution was

concentrated to give quantitative yield of a yellow syrup. The product was analysed by LC-MS (ES ionisation) giving $[MH]^+$ at 335 as expected. 1H (500 MHz) and ^{13}C (125 MHz) NMR spectroscopy was in agreement with structure

5 The product was used in the next step without further purification.

17-Amino-5-oxo-6-aza-3,9,12,15-tetraoxaheptadecanoic acid

10 A solution of 17-azido-5-oxo-6-aza-3,9,12,15-tetraoxaheptadecanoic acid (8.36 g, 25.0 mmol) in water (100 ml) was reduced using $H_2(g)$ -Pd/C (10%). The reaction was run until LC-MS analysis showed complete conversion of starting material (column Vydac 218TP54; solvents: A =

15 water/0.1% TFA and B = acetonitrile/0.1% TFA; gradient 4-16% B over 20 min; flow 1.0 ml/min; UV detection at 214 and 284 nm, ES ionisation giving $M+H$ at 335 for starting material and 309 for the product). The solution was filtered and used directly in the next step.

20

17-(Fmoc-amino)-5-oxo-6-aza-3,9,12,15-tetraoxaheptadecanoic acid


To the aqueous solution of 17-amino-5-oxo-6-aza-3,9,12,15-tetraoxaheptadecanoic acid from

25 above (corresponding to 25.0 mmol amino acid) was added sodium bicarbonate (5.04 g, 60.0 mmol) and dioxan (40 ml). A solution of Fmoc-chloride (7.11 g, 0.275 mol) in dioxan (40 ml) was added dropwise. The reaction mixture was stirred overnight. Dioxan was evaporated off

30 (rotavapor) and the aqueous phase was extracted with ethyl acetate. The aqueous phase was acidified by addition of hydrochloric acid and precipitated material was extracted into chloroform. The organic phase was dried ($MgSO_4$), filtered and concentrated to give 11.3 g

(85%) of a yellow syrup. The structure was confirmed by LC-MS analysis (column Vydac 218TP54; solvents: A = water/0.1% TFA and B = acetonitrile/0.1% TFA; gradient 40-60% B over 20 min; flow 1.0 ml/min; UV detection at 214 and 254 nm, ES ionisation giving M+H at 531 as expected for the product peak at 5,8 minutes). The analysis showed very low content of side products and the material was used without further purification.


10 1 b) Synthesis of ClCH₂CO-Lys-Cys(tBu)-Arg-Gly-Asp-Cys(tBu)-Phe-Cys-PEG-NH₂



The PEG unit was coupled manually to Rink Amide AM resin, starting on a 0.25 mmol scale, mediated by HATU activation. The remaining peptide was assembled on an ABI 433A automatic peptide synthesiser using 1 mmol amino acid cartridges. The amino acids were pre-activated using HBTU before coupling. N-terminal amine groups were chloroacetylated using a solution of chloroacetic anhydride in DMF for 30 min. The simultaneous removal of peptide and side-chain protecting groups (except tBu) from the resin was carried out in TFA containing TIS (5 %), H₂O (5 %) and phenol (2.5 %) for two hours. After work-up 322 mg of crude peptide was obtained (Analytical HPLC: Gradient, 5-50 % B over 10 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; column, Phenomenex Luna 3 μ C18 (2) 50 x 4.6 mm; flow, 2 mL/min; detection, UV 214 nm; product retention time, 6.37 min). Further product

characterisation was carried out using mass spectrometry: Expected, M+H at 1409, found, at 1415).


1 c) Synthesis of thioether cyclo[CH₂CO-Lys-Cys(tBu)-Arg-Gly-Asp-Cys(tBu)-Phe-Cys]-PEG-NH₂



322 mg of ClCH₂CO-Lys-Cys(tBu)-Arg-Gly-Asp-Cys(tBu)-Phe-Cys-(PEG)_n-NH₂ was dissolved in water/acetonitrile. The mixture was adjusted to pH 8 with ammonia solution and stirred for 16 hours. After work-up crude peptide was obtained (Analytical HPLC: Gradient, 5-50 % B over 10 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; column, Phenomenex Luna 3 μ C18 (2) 50 x 4.6 mm; flow, 2 mL/min; detection, UV 214 nm; product retention time, 6.22 min). Further product characterisation was carried out using mass spectrometry: Expected, M+H at 1373, found, at 1378).

20

1 d) Synthesis of disulphide [Cys²⁻⁶] thioether cyclo[CH₂CO-Lys-Cys²-Arg-Gly-Asp-Cys⁶-Phe-Cys]-PEG-NH₂




Thioether cyclo[CH₂CO-Lys-Cys(tBu)-Arg-Gly-Asp-Cys(tBu)-Phe-Cys]-(PEG)_n-NH₂ was treated with a solution of anisole (200 μL), DMSO (2 mL) and TFA (100 mL) for 60 min following which the TFA was removed in vacuo and the peptide precipitated by the addition of diethyl ether.

Purification by preparative HPLC (Phenomenex Luna 5 μ C18 (2) 250 x 21.20 mm column) of 70 mg crude material was carried out using 0-30 % B, where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA, over 40 min at a flow rate of 10 mL/min. After lyophilisation 46 mg of pure material was obtained (Analytical HPLC: Gradient, 0-30 % B over 10 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; column, Phenomenex Luna 3 μ C18 (2) 50 x 4.6 mm; flow, 2 mL/min; detection, UV 214 nm; product retention time, 6.80 min).

Further product characterisation was carried out using mass spectrometry: Expected, M+H at 1258.5, found, at 1258.8).

1 e) Synthesis of disulfide [Cys²⁻⁶] thioether cyclo[CH₂CO-Lys(fluorescein)-Cys²-Arg-Gly-Asp-Cys⁶-Phe-Cys]-PEG-NH₂



30 mg of [Cys²⁻⁶] cyclo[CH₂CO-Lys-Cys-Arg-Gly-Asp-Cys-Phe-Cys]-PEG-NH₂, 16.2 mg of Fluorescein NHS ester and 4 μL of

N-methylmorpholine was dissolved in DMF (3 mL). The mixture was protected against light and stirred over night. Purification by preparative HPLC (Vydac 218TP1022 C18 column) of the reaction mixture was carried out using
 5 20-30 % B, where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA, over 40 min at a flow rate of 10 mL/min. After lyophilisation 21.6 mg of pure material was obtained (Analytical HPLC: Gradient, 10-40 % B over 10 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; column,
 10 Phenomenex Luna 3 μ C18 (2) 50 x 4.6 mm; flow, 2 mL/min; detection, UV 214 nm; product retention time, 7.0 min). Further product characterisation was carried out using mass spectrometry: Expected, M+H at 1616.5, found, at 1616.3).


15

Example 2:

Synthesis of disulphide [Cys²⁻⁶]thioether cyclo[CH₂CO-Asp-Cys²-Arg-Gly-Asp-Cys-Phe-Cys-Gly]-Bis(aminoethyl)ethylene glycol-fluorescein

20


2a. Synthesis of ClCH₂CONH-Asp-Cys(tBu)-Arg-Gly-Asp-Cys(tBu)-Phe-Cys-Gly-NH-(CH₂CH₂O)₂CH₂CH₂NH₂



The peptide was synthesised on a ABI 433A automatic
 25 peptide synthesiser starting with O-Bis-(aminoethyl)ethylene glycol trityl resin on a 0.25 mmol scale using 1 mmol amino cartridges. The amino acids were pre-activated using HBTU before coupling. The N-terminal was chloroacetylated using chloroacetic anhydride. The

simultaneous removal of peptide and side-chain protecting groups (except tBu) from the resin was carried out in TFA containing TIS (5 %), H₂O (5 %) and phenol (2.5 %) for two hours. After work-up 364 mg of crude peptide was obtained
 5 (Analytical HPLC: Gradient, 5-50 % B over 10 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; 2 mL/min; column, Phenomenex Luna 3 μ C18 (2) 50 x 4.6 mm; detection, UV 214 nm; product retention time, 6.55 min). Further product characterisation was carried out using electrospray mass spectrometry: expected, M+H at 1293.5, found, at 1293.4).
 10


2b. Synthesis of cyclo[CH₂CONH-Asp-Cys(tBu)-Arg-Gly-Asp-Cys(tBu)-Phe-Cys]-Gly-NH-(CH₂CH₂O)₂CH₂CH₂NH₂



15


250 mg of ClCH₂CONH-Asp-Cys(tBu)-Arg-Gly-Asp-Cys(tBu)-Phe-Cys-Gly-NH-(CH₂CH₂O)₂CH₂CH₂NH₂ was dissolved in water/acetonitrile. The mixture was adjusted to pH 8 with ammonia solution and stirred for 18 hours.
 20 After lyophilisation the crude peptide was obtained as a white powder. (Analytical HPLC: Gradient, 5-50 % B over 10 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; flow, 2 mL/min; column, Phenomenex Luna 3 μ C18 (2) 50 x 4.6 mm; detection, UV 214 nm; product retention time, 6.17 min). Further product characterisation was carried out using electrospray mass spectrometry: expected, M+H at 1257.5, found, at 1257.6).
 25

2c. Synthesis of [Cys²⁻⁶] cyclo[CH₂CONH-Asp-Cys-Arg-Gly-Asp-Cys-Phe-Cys]-Gly-NH-(CH₂CH₂O)₂CH₂CH₂NH₂



- 5 Cyclo[CH₂CONH-Asp-Cys(tBu)-Arg-Gly-Asp-Cys(tBu)-Phe-Cys]-Gly-NH-(CH₂CH₂O)₂CH₂CH₂NH₂ was dissolved in a solution of anisole (500 µl), DMSO (4 ml) and TFA (200 ml). The mixture was stirred at room temperature for 15 min following which the TFA was removed in vacuo and the peptide precipitated by the addition of diethyl ether.
- 10 Purification by preparative HPLC (Phenomenex Luna 5µ C18 (2) 250 x 21.20 mm column) of the crude material was carried out using 0-30 % B, where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA, over 40 min at a flow rate of 10 mL/min. After lyophilisation 44 mg of pure material was obtained. (Analytical HPLC: Gradient, 0-30 % B over 10 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; flow, 2 mL/min; column, Phenomenex Luna 3µ C18 (2) 50 x 4.6 mm; detection, UV 214 nm; product retention time, 5.88 min).
- 15
- 20 Further product characterization was carried out using electrospray mass spectrometry: expected, M+H at 1143.4, found, at 1143.5).

2d. Conjugation of [Cys²⁻⁶] cyclo[CH₂CONH-Asp-Cys-Arg-Gly-Asp-Cys-Phe-Cys]-Gly-NH-(CH₂CH₂O)₂CH₂CH₂NH₂ with fluorescein




- 5 10 mg of [Cys²⁻⁶] cyclo[CH₂CONH-Asp-Cys-Arg-Gly-Asp-Cys-Phe-Cys]-Gly-NH-(CH₂CH₂O)₂CH₂CH₂NH₂, 4.7 mg of fluorescein NHS ester and 5 μ L of 4-methylmorpholine were dissolved in DMF (0.5 mL) and the solution stirred for 3 hours.
- Purification by preparative HPLC (Phenomenex Luna 5 μ C18
- 10 (2) 250 x 21.20 mm column) of the crude material was carried out using 0-30 % B, where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA, over 40 min at a flow rate of 10 mL/min. After lyophilisation 6 mg of pure material was obtained. (Analytical HPLC: Gradient, 0-30 % B over 10
- 15 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; flow, 2 mL/min; column, Phenomenex Luna 3 μ C18 (2) 50 x 4.6 mm; detection, UV 214 nm; product retention time, 10.07 min). Further product characterisation was carried out using electrospray mass spectrometry: expected, M+H at 1501.4,
- 20 found, at 1501.4).

Claims:

1. A compound comprising a peptidic vector and at least one fluorescein dye, wherein the peptidic vector
5 comprises the amino acid sequence X_3 -G-D, wherein the peptidic vector and the fluorescein dyes are coupled, and wherein
 X_3 represents arginine, N-methylarginine or an arginine mimetic,
10 G represents glycine,
D represents aspartic acid,
or a physiologically acceptable salt thereof.

2. A compound as claimed in claim 1 given by formula I,



comprising two cyclising bridges,
wherein X_3 , G and D are defined as in claim 1, and wherein

R_a represents $-(CH_2)_n-$ or $-(CH_2)_n-C_6H_4-$ forming a bridge to
20 either X_2 , X_4 or X_6 , wherein
n represents a positive integer from 1 to 10, and

X_1 represents a bond or 1, 2, 3, 4 or 5 amino acid residues, wherein one amino acid residue is optionally
25 functionalised with a spacer moiety, or said amino acid residue possesses a functional side-chain such as an acid or amine group,

X_2 and X_4 represent independently amino acid residues
30 capable of forming a cyclising bridge,

X_5 represents a hydrophobic amino acid or derivatives thereof, and

X₆ represents an amino acid residue capable of forming a cyclising bridge, and


5 X₇ represents a spacer or biomodifier moiety or is absent,

and,

the compound further comprising at least one group Z,
10 representing a fluorescein dye, linked to one or more of the groups X₁, X₆ or X₇ optionally via a spacer group.


3. A compound as claimed in claim 2 selected from one of the formulas;

15




(III)

20



(IV)

25



wherein R_a , X_1 , X_3 , G , D , X_5 , and X_7 are as defined in
 5 claims 1 and 2 and wherein

X_2 , X_4 and X_6 represent amino acid residues capable of
 forming a disulphide or a thioether bond,

10 W_1 is a spacer moiety or is absent,

h is a positive integer 1 or 2,

and wherein at least one of the Z groups is present
 15 representing a fluorescein dye.

4. A compound of formula III as claimed in claim 3
 wherein R_a represents $-(\text{CH}_2)-$.

20 5. A compound of formula III as claimed in any of
 claims 3 or 4 wherein X_1 represents an amino acid residue
 with a functional side-chain such as an acid or amine
 group, the amino acid being selected from aspartic acid,
 lysine, glutamic acid, homolysine or a diaminoalicyclic
 25 acid or derivatives thereof.

6. A compound of formula III as claimed in any of
 claims 3 to 5 wherein X_2 , X_4 and X_6 independently
 represent a cysteine or homocysteine residue.

7. A compound of formula III as claimed in any of claims 3 to 6 wherein X_3 represents arginine.
8. A compound of formula III as claimed in any of
5 claims 3 to 7 wherein X_5 represents phenylalanine, tyrosine, a 3-iodo-tyrosine or naphthylalanine.
9. A compound of formula III as claimed in any of
10 claims 3 to 8 wherein X_7 comprises 1-10 units of a monodisperse PEG building block or is absent.
10. A pharmaceutical composition comprising an effective amount of a compound of any of the previous claims, together with one or more pharmaceutically acceptable
15 adjuvants, excipients or diluents.
11. Compounds of any of claims 1 to 9 for use as an optical imaging contrast agent.
- 20 12. Use of a compound as claimed in any one of claims 1 to 9 in the manufacture of an optical imaging contrast agent for use in a method of diagnosis involving administering said contrast agent to a human or animal body and generating an image of at least part of said
25 body.
13. A method of generating images of a human or animal body by optical imaging involving administering a contrast agent to said body, and generating an image of
30 at least a part of said body to which said contrast agent has distributed, characterized in that said contrast agent comprises a compound as claimed in any one of claims 1 to 9.

SEQUENCE LISTING

<110> Amersham Health AS

5 <120> Peptide based compounds

<130> PN0353

<150> NO20033115

10 <151> 2003-06-08

<160> 8

<170> PatentIn version 3.1

15 <210> 1

<211> 8

<212> PRT

<213> Artificial sequence

20 <220>

<223> Synthesised peptide

<400> 1

25 Lys Cys Arg Gly Asp Cys Phe Cys
1 5

30 <210> 2

<211> 8

<212> PRT

<213> Artificial sequence

35 <220>

<223> Synthesised peptide

<220>

<221> THIOETH

40 <222> (1)..(8)

<223> Thioether bridge between amino acids 1 and 8.

<220>

45 <221> DISULFID

<222> (2)..(6)

<223> Disulphide bridge between amino acids 2 and 6.

<220>

50 <400> 2

Lys Cys Arg Gly Asp Cys Phe Cys
1 5

55 <210> 3

<211> 10

<212> PRT

<213> Artificial sequence

60 <220>

<223> Synthesised peptide

<400> 3

5 Lys Asp Cys Arg Gly Asp Cys Phe Cys Gly
1 5 10

<210> 4
10 <211> 10
<212> PRT
<213> Artificial sequence

<220>
15 <223> Synthesised peptide

<220>
<221> THIOETH
<222> (1)..(3)
20 <223> Thioether bridge between amino acids 1 and 3

<220>
<221> DISULFID
25 <222> (7)..(9)
<223> Disulphide bridge between amino acids 7 and 9

<400> 4

30 Lys Asp Cys Arg Gly Asp Cys Phe Cys Gly
1 5 10

35 <210> 5
<211> 8
<212> PRT
<213> Artificial sequence

40 <220>
<223> Synthesised peptide

<400> 5

45 Lys Cys Arg Gly Asp Cys Phe Cys
1 5

<210> 6
50 <211> 8
<212> PRT
<213> Artificial sequence

<220>
55 <223> Synthesised peptide

<220>
<221> THIOETH
<222> (1)..(6)
60 <223> Thioether bridge between amino acids 1 and 6

<220>
<221> DISULFID
<222> (2)..(8)
5 <223> Disulphide bridge between amino acids 2 and 8.

<400> 6

10 Lys Cys Arg Gly Asp Cys Phe Cys
1 5

<210> 7
15 <211> 8
<212> PRT
<213> Artificial sequence

<220>
20 <223> Synthesised peptide

<400> 7

Asp Cys Arg Gly Asp Cys Phe Cys
25 1 5

<210> 8
<211> 8
30 <212> PRT
<213> Artificial sequence

<220>
35 <223> Synthesised peptide

<220>
<221> THIOETH
<222> (1)..(8)
40 <223> Thioether bridge between amino acids 1 and 8

<220>
<221> DISULFID
<222> (2)..(6)
45 <223> Disulphide bridge between amino aids 2 and 6

<400> 8

50 Asp Cys Arg Gly Asp Cys Phe Cys
1 5

55

INTERNATIONAL SEARCH REPORT

International Application No

PCT/N02004/000208

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/705 C07K5/09 A61K38/17 A61K38/06 A61K49/00
G01N33/58 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, PAJ, CHEM ABS Data, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2002/102217 A1 (HELLEBUST HALLDIS ET AL) 1 August 2002 (2002-08-01)	1,10-13
Y	page 14; claims 11,19,31; example 13	2-9
X	KATADA J ET AL.: "A Novel Motif for Platelet Fibrinogen Receptor Recognition" THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 272, no. 12, 21 March 1997 (1997-03-21), pages 7720-7726, XP002303382	1,10
Y	abstract; figure 2	2-9
X	DE 198 08 591 A (UNIV LEIPZIG) 16 September 1999 (1999-09-16)	1,10
Y	column 3, line 20 - line 30	2-9
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

2 November 2004

Date of mailing of the international search report

06/12/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Schmidt, Harald

International Application No
T/NO2004/000208

...T/N02004/000208

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	RIECKE B ET AL.: "Topical Application of Integrin Antagonists Inhibits Proliferative Retinopathy" HORMONE AND METABOLIC RESEARCH, vol. 33, no. 5, May 2001 (2001-05), pages 307-311, XP008037920	1,10
Y	the entire document, particularly page 308, right-hand column; Figure 2 and abstract	2-9
Y	WO 02/26776 A (CUTHBERTSON ALAN ; NYCOMED IMAGING AS (NO)) 4 April 2002 (2002-04-04) cited in the application the entire document, particularly pages 6-9,16,17,23,25 and the claims	2-9
Y	WO 01/77145 A (INDREVOLL BAARD ; CUTHBERTSON ALAN (NO); NYCOMED IMAGING AS (NO)) 18 October 2001 (2001-10-18) cited in the application the entire document, particularly pages 7,8,27,34 and the claims	2-9
A	DE GROOT F M H ET AL: "Design, synthesis, and biological evaluation of a dual tumor-specific motive containing integrin-targeted plasmin-cleavable doxorubicin prodrug" MOLECULAR CANCER THERAPEUTICS, AMERICAN ASSOCIATION OF CANCER RESEARCH, US, vol. 1, September 2002 (2002-09), pages 901-911, XP002978656 ISSN: 1535-7163 abstract; figure 1	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/N02004/000208

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 13 is directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

T/NO2004/000208

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2002102217 A1	01-08-2002	US 6331289 B1	18-12-2001
		AT 249247 T	15-09-2003
		AU 733477 B2	17-05-2001
		AU 4718297 A	22-05-1998
		AU 733495 B2	17-05-2001
		AU 4786697 A	22-05-1998
		AU 4786797 A	22-05-1998
		AU 4786897 A	22-05-1998
		AU 4786997 A	22-05-1998
		AU 4787097 A	22-05-1998
		BG 103438 A	31-01-2000
		BG 103439 A	31-01-2000
		BR 9712683 A	19-10-1999
		BR 9713978 A	02-05-2000
		CA 2269985 A1	07-05-1998
		CA 2270120 A1	07-05-1998
		CN 1440816 A	10-09-2003
		CN 1238700 A	15-12-1999
		CN 1234742 A	10-11-1999
		CZ 9901494 A3	15-09-1999
		DE 69724817 D1	16-10-2003
		DE 69724817 T2	01-07-2004
		EP 1007101 A2	14-06-2000
		EP 0973552 A2	26-01-2000
		EP 0991427 A2	12-04-2000
		EP 0971747 A2	19-01-2000
		EP 0946202 A2	06-10-1999
		EP 0963209 A2	15-12-1999
		ES 2206689 T3	16-05-2004
		WO 9818500 A2	07-05-1998
		WO 9818501 A2	07-05-1998
		WO 9818495 A2	07-05-1998
		WO 9818496 A2	07-05-1998
		WO 9818497 A2	07-05-1998
		WO 9818498 A2	07-05-1998
		HU 0000357 A2	28-06-2000
		JP 2001511765 T	14-08-2001
		JP 2001503407 T	13-03-2001
		JP 2002515889 T	28-05-2002
		JP 2001502719 T	27-02-2001
		KR 2000052829 A	25-08-2000
		KR 2000052830 A	25-08-2000
		NO 991889 A	28-06-1999
		NO 991890 A	28-06-1999
		NZ 335596 A	27-10-2000
		NZ 335799 A	24-11-2000
		US 6264914 B1	24-07-2001
		US 6051207 A	18-04-2000
DE 19808591 A	16-09-1999	DE 19808591 A1	16-09-1999
WO 0226776 A	04-04-2002	AU 9245601 A	08-04-2002
		CA 2420577 A1	04-04-2002
		CN 1462276 T	17-12-2003
		EP 1358206 A2	05-11-2003
		JP 2004509975 T	02-04-2004
		WO 0226776 A2	04-04-2002
		NO 20031323 A	24-03-2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/N02004/000208

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0226776	A	US 2003176639 A1	18-09-2003
WO 0177145	A	18-10-2001	
		AU 5068301 A	23-10-2001
		CA 2405469 A1	18-10-2001
		CN 1436195 T	13-08-2003
		EP 1272507 A2	08-01-2003
		JP 2003531835 T	28-10-2003
		WO 0177145 A2	18-10-2001
		US 2003204049 A1	30-10-2003